

## DRUG FACTS

### Active ingredient(s)

### Purpose

#### Use(s)

- relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

#### Warnings

**Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

#### Do not use

- to prevent hives from any known cause such as:
  - foods
  - insects stings
  - medicines
  - latex or rubber gloves

because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.

- if you have ever had an allergic reaction to this product (loratadine) or any of its ingredients

#### Ask a doctor before use if you have

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

#### Ask a doctor or pharmacist before use if you are

#### When using this product

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Get medical help right away**
- symptoms don't improve after 3 days of treatment
- the hives have lasted more than 6 weeks

#### If pregnant or breast-feeding

#### Keep out of reach of children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

adults and children 6 years and over:	2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours
children under 6 years of age:	ask a doctor
consumers with liver or kidney disease:	ask a doctor

#### Other information

- Safety sealed. Do not use if imprinted seal around cap is broken or missing
- Store at room temperature

Manufactured by:  
Silarx Pharmaceuticals, Inc.  
19 West Street  
Spring Valley, NY 10977 USA

#### Inactive ingredients

citric acid, edetate disodium, flavor, glycerin, propylene glycol, sodium benzoate, sugar, water.